QL
483
I 4C3X
pt.8
Ent.
F

Catalogue of Indian Insects

PART 8—Amatidae (Syntomidae)

BY

T. BAINBRIGGE FLETCHER,

R.N., F.L.S., F.E.S., F.Z.S.,

Imperial Entomologist.

SMITHSONIAN INSTITUTION
AUG 6 1925
268486
NATIONAL MUSEUM

Issued under the authority of the Government of India.

CALCUTTA: GOVERNMENT OF INDIA
CENTRAL PUBLICATION BRANCH
1925

QL
483
I 4C3X
pt.8
Ent.
F

Catalogue of Indian Insects

PART 8—Amatidae (Syntomidae)

BY

T. BAINBRIGGE FLETCHER,

R.N., F.L.S., F.E.S., F.Z.S.,

Imperial Entomologist.

SMITHSONIAN INSTITUTION
AUG 6 1925
268486
NATIONAL MUSEUM

Issued under the authority of the Government of India.

CALCUTTA: GOVERNMENT OF INDIA
CENTRAL PUBLICATION BRANCH
1925

Government of India Publications are obtainable from the Government of India Central Publication Branch, 8, Hastings Street, Calcutta, and from the following Agents :—

EUROPE.

OFFICE OF THE HIGH COMMISSIONER FOR INDIA,
42, GROSVENOR GARDENS, LONDON, S.W. 1.

Ernest Leroux, 28, Rue Bonaparte, Paris.
Martinus Nijhoff, The Hague, Holland.
Otto Harrassowitz, Leipzig.
Friedländer und Sohn, Berlin.
and all Book-sellers in the United Kingdom.

INDIA AND CEYLON.

Provincial Book Depôts :

MADRAS :—Office of the Superintendent, Government Press, Mount Road, Madras.
BOMBAY :—Office of the Manager, Government Central Press, Bombay.
SIND :—Library attached to the Office of the Commissioner in Sind, Karachi.
BENGAL :—Office of the Bengal Secretariat Book Depôt, Writers' Buildings, Room No. 1, Ground Floor, Calcutta.
UNITED PROVINCES OF AGRA AND OUDH :—Office of the Superintendent of Government Press, United Provinces of Agra and Oudh, Allahabad.
PUNJAB :—Office of the Government of Punjab, Lahore.
BURMA :—Office of the Superintendent, Government Printing, Burma, Rangoon.
CENTRAL PROVINCES AND BERAR :—Office of the Central Provinces Secretariat, Nagpur.
ASSAM :—Office of the Superintendent, Assam Secretariat Press.
BIHAR AND ORISSA :—Office of the Superintendent, Government Printing, Bihar and Orissa, P. O. Gulzarbagh, Patna.
COORG :—Office of the Chief Commissioner of Coorg, Bangalore.
NORTH-WEST FRONTIER PROVINCE :—Government Press, Peshawar.

Thacker, Spink & Co., Calcutta and Simla.
W. Newman & Co., Ltd., Calcutta.
R. Cambray & Co., Calcutta.
S. K. Lahiri & Co., Calcutta.
The Indian School Supply Depôt, 309, Bow Bazar Street, Calcutta, and 226, Nawabpur, Dacca.
Butterworth & Co. (India), Ltd., Calcutta.
Rai M. C. Sarcar Bahadur & Sons, 90-2A, Harrison Road, Calcutta.
The Weldon Library, 17, Park Street, Calcutta.
Standard Literature Company, Limited, Calcutta.
Association Press, Calcutta.
Chukervertty, Chatterjee & Co., Ltd., 13, College Square, Calcutta.
Higginbotham & Co., Madras.
V. Kalyanarama Iyer & Co., Madras.
P. R. Rama Iyer & Co., Madras.
Rochouse and Sons, Madras.
Bright & Co., Trivandrum.
V. S. Swaminathan, Bookseller, West Tower Street, Madura.
Thacker & Co., Ltd., Bombay.
D. B. Taraporevala, Sons & Co., Bombay.
Sunder Pandurang, Bombay.
Ram Chandra Govind & Sons, Kalbadevi, Bombay.
N. M. Tripathi & Co., Booksellers, Princess Street, Kalbadevi Road, Bombay.
Proprietor, New Kitabkhana, Poona.
The Manager, Oriental Book Supplying Agency, 15, Shukrawar, Poona City.
R. S. Gondhalekar's Book Depôt, Publisher and Bookseller, Budhwar Chawk, Poona City.
Managing Director, Co-operative Bookstall, Booksellers and Publishers, Poona City.
The Standard Bookstall, Karachi, Rawalpindi, Murree, Lahore, Peshawar and Quetta.
The Book Company, Calcutta.
Burma Book Club Ltd., Rangoon
Karsandas Narandas & Sons, Surat.
A. H. Wheeler & Co., Allahabad, Calcutta and Bombay.
N. B. Mathur, Supdt., Nazir Kanun Hind Press, Allahabad.
The North India Christian Tract and Book Society, 18, Clive Road, Allahabad.
Ram Dayal Agarwala, 184, Katra, Allahabad.
Manager, Newal Kishore Press, Lucknow.
The Upper India Publishing House, Ltd., 41, Aminabad Park, Lucknow.
Munshi Seeta Ram, Managing Proprietor, Indian Army Book Depôt, Juhi, Cawnpore.
Rai Sahib M. Gulab Singh & Sons, Mufid-i-Am Press, Lahore and Allahabad.
Rama Krishna & Sons, Booksellers, Anarkall, Lahore.
Puri Brothers, Booksellers and Publishers, Katcheri Road, Lahore.
Manager of the Imperial Book Depôt, 63, Chandney Chawk Street, Delhi.
Oxford Book and Stationery Company, Delhi.
Supdt., American Baptist Mission Press, Rangoon.
Proprietor, Rangoon Times Press, Rangoon.
The Modern Publishing House, Ltd., 30, Phayre Street, Rangoon.
The International Buddhist Book Depôt, Post Box No. 971, Rangoon.
Manager, the "Hitavada," Nagpur.
S. C. Talukdar, Proprietor, Students & Co., Cooch Behar.
Times of Ceylon Co., Ltd.
The Manager, Ceylon Observer, Colombo.
The Manager, The Indian Book Shop, Benares City.
B. C. Basak, Esq., Proprietor, Albert Library, Dacca.

Catalogue of Indian Insects.

AMATIDÆ (SYNTOMIDÆ).

Syntomidæ: Hampson, F. I. Moths I 209-228 (1892) [key to genera]; *id.*, Cat. Lep. Phal. I (1898) [revision; keys to genera and species]; Seitz, Lep. Pal. II 37-42 (1910); *id.*, Lep. Indo-Austr. II 61-92 (1912-1913).

Amatidæ: Hampson, Cat. Phal. Suppl. I, pp. 1-388 (1914).

1. Psichotoë, Boisduval (1829).

Psichotoë, Boisd., Mon. Zyg., p. 129 (1829); Hampson, F. I. Moths I 225, f. 141 (1892) [redescr.]; *id.*, Cat. Phal. I 32, f. 9 (1898) [redescr.]; Seitz, Lep. Indo-Austr., II 63 (1912). Type: *duvauceli*, Boisd.

Cacosoma, Felder, Reise Novara, Lep. IV, p. 2 (1874). Type *gnatula*, Boisd. [S. Africa].

duvauceli, Boisduval—

Psichotoë duvauceli, Boisd., Mon. Zyg., p. 129, t. 8, f. 5 (1829)[1]; Hmpsn., F. I. Moths I 226, f. 141 (1892) [redescr.][2]; *id.*, Cat. Phal. I 32, f. 9 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 63, t. 10, f. A (1912).[4]

Karachi. [2, 4]
Lahore.
Pusa.
Calcutta. [2, 3]
Gauhati.
Jorhat.
Assam. [3, 4]
Burma. [3, 4]
Rangoon.
Minbu.
Mandalay.
Hsipaw.
Pyinmana.
Pegu.
Mergui.

2. Callitomis, Butler (1876).

Callitomis, Butl., J. Linn. Soc. XII 351, t. 27, f. 1 (1876); Hmpsn., F. I. Moths I 224, f. 139 (1892) [redescr.]; *id.*, Cat. Phal. I 33 (1898) [redescr.]; *id.*, B. J. XIII 45 (1900) [key to spp.]; Seitz, Lep. Pal. II 41 (1910) [redescr.]; *id.*, Lep. Indo-Austr. II 63 (1912) [redescr.]. Type: *syntomoides*, Butl.

leucosoma, Butler—

Callitomis leucosoma, Butl., J. Linn. Soc. XII 351 (1876)[1]; Hmpsn., F. I. Moths I 225 (1892) [redescr.][2]; *id.*, Cat. Phal. I 34, t. 1, f. 3 (1898) [redescr.][3]; Seitz, Lep. Pal. II 41, t. 9, f. H (1910).[4] — Kashmir. [1-4]

multifasciata, Hampson.

Callitomis multifasciata, Hmpsn., F. I. Moths I 225 (1892)[1]; *id.*, Cat. Phal. I 33, t. 1, f. 1 (1898) [redescr.][2]; Seitz, Lep. Indo-Austr. II 63, t. 10, f. A (1912)[3]. — Nagas. [1-3] Sikkim. [2, 3]

syntomoides, Butler.

Callitomis syntomoides, Butl., J. Linn. Soc. XII 351 (1876)[1]; Hmpsn., F. I. Moths I 224, f. 139 (1892) [redescr.][2]; *id.*, Cat. Phal. I 34, f. 12 (1898) [redescr.][3]; Seitz, Lep. Pal. II 41 (1910)[4]; *id.*, Lep. Indo-Austr. II 63 (1912).[5] — Kashmir. [1-4] Murree. [3] Simla. [2]

3. Ceryx, Wallengren (1863).

Ceryx, Wlgn., Wien. Ent. Mon. VII 140 (1863); Hmpsn., Cat. Phal I 35 (1898) [redescr.]; *id.*, B. J. XIII 45-46 (1900) [key to spp.]; Seitz, Lep. Pal. II 41 (1910) [redescr.; habits]; *id.*, Lep. Indo-Austr. II 63 (1912) [redescr.]. Type: *anthraciformis*, Wlgn. [S. Africa].

Agaphthora Meyrick, Proc. Linn. Soc. N. S. Wales (2) I 774 (1886). Type: *guttulosa*, Wlk. [Papuan subregion].

Syntomoides, Hmpsn., F. I. Moths I 209-210, f. 137 (1892). Type: *imaon* Cram.

cherra, Moore.

Syntomis cherra, Mo., Lep. Atk., pp. 12-13 (1879)[1]; Hmpsn., F. I. Moths I 220 (1892) [redescr.][2].	Khasis. [1-8] Shillong. [3], [4] Cherrapunji. [1]
Syntomoides quisqualis, Swinh., A.M.N.H. (6) XIV 441 (1894)[3]; *id.*, T. E. S. 1895, 30 (1895)[4]; Hmpsn., F. I. Moths IV 465 (1896).[5]	
Ceryx cherra, Hmpsn., Cat. Phal. I 41, t. 1, f. 13 (1898) [redescr.][6]; *id.*, B. J. XIII 46 (1900)[7]; Seitz, Lep. Indo-Austr. II 65 t. 10, f. D (1912).[8]	

diptera, Fabricius.

Zygæna diptera Fb., Syst. Ent. p. 555 (1775).[1]	S. India. [4], [11] Tranquebar.
Sphinx atereus, Stoll, Pap. Exot. IV 240, t. 400, f. A (1782)[2].	Coromandel Coast. [2], [7] Coimbatore. [3], [8]
Syntomis incipiens, Wlk., Cat. XXXI 68-69 (1864).[3]	
Syntomoides incipiens, Hmpsn., F. I. Moths I 211 (1892).[4]	
Syntomoides brachypecten, Hmpsn., F. I. Moths I 211 (1892)[5].	Ceylon. [8], [11]
Psychotoë brachypecten, Hmpsn., Ill. Het. IX 61, t. 157, f. 23 (1893).[6]	Kanthalai. [4]
Syntomis aterea, Hmpsn., F. I. Moths I 223 (1892).[7]	Anuradhapura. [5]
Ceryx diptera, Hmpsn., Cat. Phal. I 36-37, f. 15 (1898) [redescr.][8]; *id.*, B. J. XIII 45 (1900)[9]; Seitz, Lep. Indo-Austr. II 64, t. 9, f. F (1912)[10].	
Ceryx diptera f. *brachypecten*, Seitz, Lep. Indo-Austr. II 64 (1912).[11]	

ginorea, Swinhoe.

Eressa ginorea, Swinh., A.M.N.H. (6) XIV 441 (1894)[1]; *id.*, T.E.S. 1895, 32, t. 1, f. 8 (1895)[2]; Hmpsn., F. I. Moths IV 465 (1896).[3]	Khasis. [1-5] Cherrapunji. [1], [2]

Ceryx ginorea, Hmpsn., Cat. Phal. I 37 (1898) [redescr.][4] ; Seitz., Lep. Indo-Austr. II 64, t. 10, f. B (1912).[5]	

codarti, Boisduval.

Syntomis godarti, Boisd., Mon. Zyg., p. 115, t. 7, f. 3 (1829)[1].	Khatmandu (Nepal). Sikkim.[6]
Syntomis libera, Wlk., Cat. XXXI 78-79 (1864).[2]	Gauhati. Shillong.
Syntomis fytchei, Mo. P.Z.S. 1871, 246, t. 18, f. 3 (1871).[3]	Cherrapunji.[9] Yunnan.[3, 6]
Syntomis cupreipennis, Butl., J. Linn. Soc. XII 347-348 (1876).[4]	Burma.[6] Lashio (U. Burma).
Syntomoides godartii, Hmpsn., F. I. Moths I 210 (1892) [*partim*].[5]	Dacca. Pusa.
Ceryx godarti, Hmpsu., Cat. Phal. I 43-44 (1898) [redescr.][6] ; *id.*, B. J. XIII 46 (1900)[7] ; Seitz, Lep. Indo-Austr. II 88, t. 9, f. F (1913)[8].	Calcutta.[9, 6] S. India.[6, 8] Ceylon.[6, 8]
Syntomoides cupreipennis, Swinh., T.E.S. 1895 30 (1895).[9]	S. China.[8] Malacca.[2, 6] Sumatra.[6]

hyalina, Moore.

Syntomis hyalina, Mo., Lep. Atk. p. 13 (1879).[1]	Sikkim.[3, 6, 7] Darjiling.[1]
Syntomis volans, Swinh., T.E.S. 1890, 173, t. 6, f. 6 (1890).[2]	
Syntomoides hyalina, Hmpsn., F. I. Moths I 211 (1892).[3]	Assam.[7]
Syntomoides volans, Hmpsn., F. I. Moths I 211 (1892).[4]	Nagas.[3, 6] Cherrapunji.
Ceryx hyalina, Hmpsn., B. J. XIII 46 (1900)[5] ; *id.*, Cat. Phal. I 42 (1898) [redescr.][6] ; Seitz, Lep. Indo-Austr. II 66, t. 9, f. G (1912).[7]	Shillong. Burma.[7] Karen Hills.[2, 4, 6]

imaon, Cramer.

Sphinx imaon, Cram., Pap. Exot. III 94, t. 248, f. E (1780).[1]	Kashmir.[12, 13] Kangra.[9]
Syntomis imaon, Wlk., Cat. I 127 (1854).[2]	Sikkim.[9] Darjiling.

Syntomis fusiformis, Wlk., Cat. VII 1595 (1856).[3] [Hindostan].	Khasis.[9] Shillong.[14, 15]
Syntomis approximata, Wlk., Cat. XXXI 79 (1864).[4]	Cachar.[6, 9]
Syntomis artina, Butl., J. Linn. Soc. XII 347 (1876)[5]; Mo., Lep. Ceylon II 35, t. 94, f. 4, (1883).[5a]	
Syntomis sargania, Butl., T.E.S. 1879 4 (1879).[6]	Calcutta.[5, 9]
Syntomis mota, Swinh., T.E.S. 1891. 134 (1891).[7]	Bombay.[9]
Syntomoides imaon, Hmpsn., F. I. Moths I 210, f. 137 (1892) [redescr.].[8]	Nilgiris.[7, 9, 13]
Ceryx imaon, Hmpsn., Cat. Phal. I 44, f. 18 (1898) [redescr.][9]; *id.*, B. J. XIII 46 (1900)[10]; Piepers, Tijds. Ent. XLVII 49[11]; Seitz, Lep. Pal. II 41, t. 9, f. H (1910)[12]; *id.*, Lep. Indo-Austr. II 65, t. 10, f. E (1912).[13]	Ceylon.[5a, 9] Kandy.[13] Lashio (U. Burma) Karen Hills.[9] Maymyo. Tenasserim.[9]
Syntomoides sargania, Swinh., T.E.S. 1895. 30 (1895).[14]	
Syntomoides fusiformis, Swinh. T.E.S. 1895. 30 (1895) [*approximata* is synon.].[15]	Hongkong.[9, 13] Sumatra.[4]

Larva descr.[11]

pleurasticta, Hampson.

Ceryx pleurasticta, Hmpsn., A.M.N.H. (7) VIII 165 (1901) [*nec pleurosticta*, Hmp. 1898][1]; *id.*, B. J. XVI 151, t. D, f. 2 (1904)[2]; *id.*, Cat. Phal. Suppl. I 6, t. 1, f. 7 (1914).[3]	N. Kanara.[1, 5]
Ceryx laterimacula, Zerny, Junk's Cat. Lep., pt. 7, p. 84; Seitz, Lep. Indo-Austr. II 88 (1913).[5]	

semicincta, Hampson.

Syntomoides semicincta, Hmpsn., T.E.S. 1895 282 (1895)[1]; *id.*, F. I. Moths IV 465 (1896) [redescr.].[2]	Tenasserim.[1, 2, 4] Dawna Hills.[3]

Ceryx semicincta, Hmpsn, Cat. Phal. I 43, t. 1, f. 16 (1898) [redescr.][3] ; Seitz, Lep. Indo-Austr. II 66, t. 10, f. E (1912)[4].

Malacca.[3], [4]
Borneo.[3], [4]

4. Trichæta, Swinhoe (1892).

Trichæta, Swinh., Cat. Het. Mus. Oxon. I 52 (1892) ; Hmpsn., F. I. Moths I 225, f. 140 (1892) [redescr.] ; *id.*, Cat. Phal. I 53 (1898) [redescr.] ; *id.*, B. J. XIII 46 (1900) [key to spp.] ; Seitz, Lep. Indo-Austr. II 66 (1912) [redescr.]. Type : *teneiformis*, Wlk.

teneiformis, Walker.

Syntomis teneiformis, Wlk., Cat. VII 1595-1596 (1856).[1] [Hindostan].
Trichæta teneiformis, Swinh., Cat. Het. Mus. Oxon. I 52 (1892)[2] ; Hmpsn., F. I. Moths I 225, f. 140 (1892) [redescr.][3] ; *id.*, Cat. Phal. I 58, f. 29 (1898) [redscr.[4]] ; Seitz, Lep. Indo-Austr. II 89 (1913)[5] ; Hmpsn., Cat. Phal. Suppl. I, p. 12 (1914).[6]
Eressa simplex, Rothschild, Nov. Zool. XVII 439 (1910)[7] ; *id.*, Nov. Zool. XIX 376, t. 5, f. 20 (1912)[8] ; Seitz, Lep. Indo-Austr. II 85 (1913) ♂[9].

Simla.[3], [4], [5], [7]
Sabathu.[9]
Bhutan.[4], [5]

5. Amata, Fabricius (1807).

Amata, Fabr., Illiger's Mag. VI 289 (1807) ; Hmpsn., Cat. Phal. Suppl. I 13 (1914). Type : *passalis*, Fb.

Syntomis, Ochsenheimer, Eur. Schmett. II 103 (1808) ; Hübner, Verz. p. 121 (1820) ; Hmpsn., F. I. Moths I 212 (1892) [redescr.] ; *id.*, Cat. Phal. I 59 (1898) [redescr.] ; *id.*, B. J. XIII 46-49 (1900) [key to spp.] ; Seitz, Lep. Pal. II 38 (1910) [redescr.] ; *id.*, Lep. Indo-Austr. II 67-68 (1912) (redescr.]. Type : *phegea*, Linn. [Europe.]

Cænochromia, Hübner, Verz. p. 121 (1820). Type : *passalis*, Fb.

Hydrusa, Walker, Cat. I 255 (1854). Type : *bicolor*, Wlk. [Queensland].

Asinusca, Wallengren, Œfv. Vet. Akad. Förh., XIX 197 (1862). Type : *atricornis*, Wlgn. [S. Africa].

Asinutea, Wlgn., Wien. Ent. Mon. VII 139 (1863). Type: *atricornis* Wlgn. [S. Africa].

Buthysia, Wlgn., Wien. Ent. Mon. VII 139 (1863). Type: *huebneri*, Boisd.

actea, Swinhoe (Two subspecies).

(1) **actea actea, Swinhoe.**

Hydrusa actea, Swinh., T.E.S. 1891. 474, t. 19, f. 7 (1891).[1] — Khasis.[1, 2, 5, 6]

Syntomis baicea [*nec* Swinh.], Hmpsn., F. I. Moths I 222 (1892) [*partim*].[2] — Shillong.[3]

Eressa actea, Swinh., T.E.S. 1895. 32 (1895).[3]

Syntomis actea, Hmpsn., B. J. XI 284 (1897) [*baicea* is distinct][4]; *id.*, Cat. Phal. I 64 (1898) [redescr.][5]; Seitz, Lep. Indo-Austr. II 74 (1913).[6]

(2) **actea swinhoei, Leech.**

Syntomis swinhoei, Leech, Entom. XXXI, 152 (1898)[1]; *id.*, T.E.S. 1898. 322 (1898).[2] — Moupin [2, 3] Chia-ting-fu.[2]

Syntomis actea subsp. 1, Hmpsn., Cat. Phal. I 64 (1898).[3] — W. China.[2, 5, 6] Chusan.[3]

Syntomis actea swinhoei, Hmpsn., Cat. Phal. I 537 (1898)[4]; Seitz, Lep. Pal. II 40 (1910)[5]; *id.*, Lep. Indo-Austr. II 74 (1913).[6]

Syntomis swinhoei var. *obsoleta*, Leech, Entom. XXXI, 152 (1898).[7]

albapex, Hampson.

Syntomis albapex, Hmpsn., F. I. Moths I 214 (1892)[1]; *id.*, Ill. Het. IX 62, t. 157, f. 1 (1893)[2]; *id.*, Cat. Phal. I 105-106 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 69, t. 9, f. G (1913).[4] — Pundaluoya.[3, 4] Nawalapitiya.[3, 4]

albifrons, Moore.

Syntomis albifrons, Mo., P.Z.S. 1878. 845, t. 53, f. 6 (1878)[1]; Hmpsn., F. I. Moths I 216 (1892) [redescr.][2]; *id.*, Cat. Phal. I 100 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 73, t. 9 f. D (1913).[4]

Tenasserim.[3] Momeit.[2] Dawna Hills.[3]

assamica, Rothschild.

Amata khasiana [*nec* Butl.], Roths., Nov. Zool. XVII 435 (1910).[1]
Amata assamica, Roths., Nov. Zool. XIX 375, t. 3, f. 21 (1912)[2]; Hmpsn., Cat. Phal. Suppl. I 31-32 (1914).[3]
Syntomis lucina assamensis, Zerny; Seitz, Lep. Indo-Austr. II 69 (1913).[4]

Khasis.[1-4]

aurea, Swinhoe.

Syntomis aurea, Swinh., A.M.N.H. (8) I. 62-63 (1908) ♂♀[1]; Seitz, Lep. Indo-Austr. II 71 (1913).[2]

Khasis.[1, 2]

aureola, Swinhoe.

Syntomis aureola, Swinh., A.M.N.H. (7) IX. 78-79 (1902) ♂♀[1]; Seitz, Lep. Indo-Austr. II 71, t. 9, f. G. (1913).[2]

Khasis.[1, 2]

baiæa, Swinhoe.

Hydrusa baiæa, Swinh., T.E.S. 1891. 473, t. 19 f. 10 (1891).[1]
Syntomis baiæa, Hmpsn., F. I. Moths I 222 (1892) [*partim*][2]; *id.*, B. J. XI 284 (1897)[3]; *id.*, Cat. Phal. I 67-68 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 74, t. 9, f. F (1913).[5]
Eressa baiæa, Swinh., T.E.S. 1895. 32 (1895)[6]

Khasis.[1, 2, 4, 5] Shillong.[6] Cherrapunji.[6]

———

Java.[4, 5] Borneo.[4, 5]

berinda, Moore.

Syntomis berinda, Mo., P.Z.S. 1878. 845, t. 53, f. 8 (1878)[1]; Hmpsn., F. I. Moths I 219 (1892) [redescr.][2]; *id.*, Cat. Phal. I 98 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. H (1913).[4]

Dacca.
Tenasserim.[1-4]
Moulmein.[3, 4]

bicincta, Kollar.

Syntomis bicincta, Kollar, Hügel's Kaschmir IV 460, t. 19, f. 8 (1844)[1]; Hmpsn., F. I. Moths I 213 (1892) [redescr.][2]; Hmpsn., Cat. Phal. I 106-107 (1898) [redescr.][3]; Seitz, Lep. Pal. II 38, t. 9, f. B (1910)[4]; *id.*, Lep. Indo-Austr. II 68 (1912).[5]

Kashmir.[4, 5]
Simla.
Kangra.[2, 3]
Kulu.[2]
Masuri.
Dehra Dun.[3]
Sikkim.[2, 3]
Calcutta.[3, 4]
Khasis.[2, 3]

Fuchau (China).[3]

chlorocera, Hampson.

Syntomis chlorocera, Hmpsn., F. I. Moths I 218 (1892)[1]; *id.*, Cat. Phal. I 93, t. 3, f. 22 (1898) [redescr.][2]; Seitz, Lep. Indo-Austr. II 74, t. 11, f. C (1913).[3]

Nagas.[1-3]

compta, Walker.

Syntomis compta, Wlk., Char. Undescr. Het. p. 89 (1869)[1]; Hmpsn., Cat. Phal. I 99, t. 5, f. 1 (1898) [redescr.][2]; Dudgeon, B. J. XII 293, t. 1, f. 24 (1899)[3]; Hmpsn., B. J. XIII 50 (1900) [redescr.][4]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. K (1913).[5]

* Benares.[1]
Khasis.[2, 4, 5]
Sikkim.[4]

* Walker described this and many other species as "collected near Benares by the late Colonel John Graham, of Her Majesty's Indian Army." It is, however, obvious that the locality given is incorrect for *S. compta* and many of the other species described at the same time. T. B. F.

confluens, Leech.

Syntomis acrospila [*nec* Feld.], Leech, T.E.S. 1898. 322 (1898)[1] ; Hmpsn., Cat. Phal. I 91-92 (1898) [*partim*].[2] — Ta-tsien-lu.[1, 2] Nitou.[1, 2] Washan.[1, 2, 7]

Syntomis acrospila var. *confluens*, Leech, T.E.S. 1898. 322 (1898).[3] — Chia-kou-ho.[1]

Amata confluens, Hmpsn., Cat. Phal. Suppl. I 33 (1914).[4] — Huang-mu-chang.[2]

Amata leechi, Roths., Novit. Zool. XVII 433 (1910)[5] ; *id.*, Novit. Zool. XIX 375, t. 3, f. 23 (1912).[6]

Syntomis leechi, Seitz, Lep. Pal. II 445 (1912).[7]

congenita, Hampson.

Amata congenita, Hmp., Nov. Zool. XXV, 93 (1918) ♂ ♀.[1] — Nilgiris.[1] Shevaroys.[1]

connectens, Rothschild.

Amata connectens, Roths., Novit. Zool. XVII 433 (1910)[1] ; *id.*, Novit. Zool. XIX 375, t. 3, f. 37 (1912)[2] ; Hmpsn., Cat. Phal. Suppl. I 33 (1914)[3] ; Seitz, Lep. Indo-Austr. II 71, t. 10, f. L (1913) [*Syntomis*].[4] — Bernardmyo.[1, 5]

Syntomis sladeni [*nec* Moore] Hmpsn., F. I. Moths I 217 (1892) [*partim*].[5]

consequa, Leech.

Syntomis consequa, Leech, Entom. XXXI 153 (1898)[1] ; Hmpsn., Cat. Phal. I 96-97, t. 5, f. 2 (1898) [redescr.][2] ; Leech, T.E.S. 1898. 321 (1898)[3] ; Seitz, Lep. Pal. II 40, t. 9, f. E (1910).[4] — Moupin.[1-3] Tibet.[4] W. China.

cyssea, Stoll.

Sphinx cysseus, Stoll, Pap. Exot. IV, t. 355, f. B (1782).[1]	Cherat (N.-W. F. P.). Kashmir.[15, 16]
Zygæna collaris, Fabr., Ent. Syst. III (*i*) p. 388 (1793).[2]	Kangra.[12, 14]
Syntomis schocnerrhi, Boisd., Mon. Zyg. p. 112, t. 7, f. 1 (1829).[3]	Nepal.[14] Darjiling.
Syntomis cyssca, Boisd., Mon. Zyg. p. 114, t. 7, f. 2 (1829).[4]	Sikkim.[12, 14] Kurseong.
Syntomis cuprea, Prittw., Stett. ent. Ztg. XXVIII, 277 (1867).[5]	Shillong.[13] Pusa.
Syntomis georgina, Butl., J. Linn. Soc. XII 345 (1876)[6]; Mo., Lep. Ceylon II 34-35, t. 95, f. 2 (1882)[7]; Hmpsn., F. I. Moths I 213 (1892) [redescr.][8]; *id.*, F. I. Moths IV 465 (1896).[9]	Calcutta.[14, 16] Karachi.[14, 16] Surat. Mhow.[14] Bombay.
Syntomis cysseoides, Butl., J. Linn. Soc. XII 346 (1876)[10]; Mo., Lep. Ceylon II 35, t. 95, f. 1 (1882).[11]	Nilgiris.[14] Coimbatore. Coorg.
Syntomis cyssea, Hmpsn., F. I. Moths I 213, f. 138 (1892) [redescr.][12]; Swinh., T.E.S. 1895. 30 (1895)[13]; Hmpsn., Cat. Phal. I 105, f. 36 (1898) [redescr.][14]; Seitz, Lep. Pal. II 39, t. 9, f. E (1910)[15]; *id.*, Lep. Indo-Austr. II 68 (1912).[16]	Ceylon.[6, 7, 10, 11, 14, 16] Peradeniya.

diaphana, Kollar.

Syntomis diaphana, Kollar, Hügel's Kaschmir IV 460, t. 19, f. 7 (1844).[1]	Kashmir.[4, 6, 7] Kasauli.[6]
Syntomis vitreata, Herr. Schäff., Aussereur. Schmett. f. 267 (1855).[2]	Simla. Dalhousie.[6]
Syntomis œnone, Butl., J. Linn. Soc. XII 344 (1876).[3]	Almora.[6] Jeolikote.
Syntomis diaphana, Hmpsn., F. I. Moths I 216 (1892)[4]; Swinh., T.E.S. 1895. 31 (1895)[5]; Hmpsn., Cat. Phal. I 67 (1898) [redescr.][6]; Seitz, Lep. Pal. II 40 (1910)[7]; *id.*, Lep. Indo-Austr. II 74, t. 9, f. G (1913).[8]	Masuri.[1, 6] Darjiling district. Assam.[8] Khasis.[6] Shillong.[5] Cherrapunji.[5] Sibsagar.[4, 6] Burma.[4, 6, 7, 8]

dichotoma, Leech.

Syntomis dichotoma, Leech, Entom. XXXI 153 (1898)[1]; *id.*, T.E.S. 1898. 323 (1898)[2]; Hmpsn., Cat. Phal. I 100, t. 4, f.1 (1898) [redescr.][3]; Seitz, Lep. Pal. II 39, t. 9, f. E (1910).[4]

Moupin (Tibet).[1-3]

W. China.[2, 4]

Syntomis dichotoma var. *concurrens*, Leech, Entom. XXXI, 153 (1898).[5]

divisa, Walker.

Syntomis divisa, Wlk., Cat. I 131 (1854)[1]; Butl., Ill. Het. I 16, t. 6, f. 14 (1877)[2]; Hmpsn., F. I. Moths I 215 (1892) [redescr.][3]; Swinh., T.E.S. 1895. 30 (1895)[4]; Hmpsn., Cat. Phal. I 98-99 (1898) [redescr.][5]; Seitz., Lep. Indo-Austr. II 70, t. 9 f. D (1913).[6]

Sikkim.[3, 5, 6]
Gauhati.
Sylhet.[1, 3, 5]
Khasis.[3, 4, 5]
Shillong.
Chin Hills.[5]
Maymyo.
Upper Tenasserim.[5, 7]

Syntomis disrupta, Mo., P.Z.S. 1878. 845, t. 53, f. 5 (1878).[7]

Dawna Hills.[5]

elongata, Hampson.

Syntomis elongata, Hmpsn., B. J. XIII 50-51 (1900)[1]; *id.*, Cat. Phal. Suppl. I 38, t. 2, f. 29 (1914) [redescr.][2]; Seitz, Lep. Indo-Austr. II 78 (1913).[3]

Belgaum.[2]
Kanara.[1-3]
Pirmâd.[1, 2]

elwesi, Rothschild.

Syntomis tigrina [*nec* Wlk.], Hmpsn., F. I. Moths I 218 (1892) ♂.[1]

Burma.[1, 4, 5]

Amata elwesi, Roths., Novit. Zool. XVII 432 (1910)[2]; *id.*, Novit. Zool. XIX 375, t. 3, f. 34 (1912).[3]

Amata elwesi, Hmpsn., Cat. Phal. Suppl. I 29 (1914) [possibly *tigrina* ♂].[4]

Syntomis elwesi, Seitz, Lep. Indo-Austr. II 71 (1913).[5]

era, Swinhoe.

Hydrusa era, Swinh., T.E.S. 1891. 473, t. 19, f. 15 (1891) ♂.[1]	Khasis.[1-4]
Syntomis era, Hmpsn., F. I. Moths I 222 (1892) [redescr.] ♀[2]; *id.*, Cat. Phal. I 68 (1898) ♂[3]; Seitz, Lep. Indo-Austr. II 80 (1913)[4].	Shillong.[5]
Eressa era, Swinh., T.E.S. 1895. 32 (1895).[5]	

euryzona, Leech.

Syntomis euryzona, Leech, Entom. XXXI 153 (1898) ♂[1]; Hmpsn., Cat. Phal. I 94, t. 3, f. 19 (1898) ♂ [redescr.][2]; Seitz, Lep. Pal. II 40, t. 9. f. G (1910).[3]	Moupin.[1, 2] Tibet.[3] ——— W. China.[3]

extensa, Walker.

Syntomis extensa, Wlk., Cat. XXXV 1863 (1866)[1]; Hmpsn., Ill. Het. VIII 3, t. 143, f. 14 (1890)[2]; *id.*, F. I. Moths I 214 (1892) [redescr.][3]; *id.*, Cat. Phal. I 103 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 72, t. 11, f. A (1913).[5]	Mahableshwar.[3, 4] Matheran.[3-5] Nilgiris.[2-5] Ootacamund.

fervida, Walker.

Syntomis fervida, Wlk., Cat. I 131 (1854)[1]; Butl., Ill. Het. I 18, t. 6, f. 16 (1877)[2]; Hmpsn., F. I. Moths I 217 (1892) ♀[3]; *id.*, Cat. Phal. I 93 (1898) ♀[4]; Seitz, Lep. Indo-Austr. II 71, t. 9, f. D (1913).[5]	Moulmein.[1, 3, 4] E. Pegu.[3, 4]

flavifrons, Hampson.

Syntomis flavifrons, Hmpsn., F. I. Moths I 215 (1892)[1]; *id.*, Cat. Phal. I 101, t. 4, f. 3 (1898) [redescr.][2]; Seitz, Lep. Indo-Austr. II 69, t. 10, f. H (1913)[3]; Swinh., T.E.S. 1895. 30 (1895).[4]	Khasis.[2] Shillong.[4] Momeit (Burma).[1, 2] ——— Indo-China.[3]

flaviguttata, Hampson.

Syntomis flaviguttata, Hmpsn., B. J. XIII 50, t. B, f. 2 (1900)[1]; Seitz., Lep. Indo-Austr. II 77 (1913)[2]; Hmpsn., Cat. Phal. Suppl. I 32 (1914) [redescr.][3] — Chitral.[1-3]

flavolavata, Rothschild.

Amata flavolavata, Roths., Novit. Zool. XVII 434 (1910)[1]; *id.*, Novit. Zool. XIX 375, t. 3, f. 24 (1912)[2]; Hmpsn., Cat. Phal. Suppl. I 33 (1914) [redescr.; may be a var. of *melas*][3]; Seitz, Lep. Indo-Austr. II 73 (1913) [*Syntomis*.][4] — Khasis.[3, 4]

formosæ, Butler.

Syntomis formosæ, Butl., J. Linn. Soc. XII 346 (1876)[1]; Hmpsn., Cat. Phal. I 98, t. 3, f. 26 (1898) [redescr.][2]; Seitz, Lep. Pal. II 39, t. 9, f. E (1910)[3]; *id.*, Lep. Indo-Austr. II 69 (1913).[4] — Margarita.[2] E. Pegu.[2] ——— Fuchau.[2, 3, 5] Chusan.[2] Formosa.[1-4]

Syntomis emma, Butl., J. Linn. Soc. XII 350-351 (1876).[5]

gelatina, Hampson.

Syntomis gelatina, Hmpsn., Ill. Het. VIII 43, t. 139, f. 1 (1890)[1]; *id.*, F. I. Moths I 214 (1892) [redescr.][2]; *id.*, Cat. Phal. I 103 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 72, t. 10, f. L (1913).[4] — Nilgiris.[1-4]

grotei, Moore.

Syntomis grotei, Mo., P.Z.S. 1871. 245, t. 18, f. 4 (1871)[1]; Hmpsn., F. I. Moths I 218 (1892) [redescr.][2]; *id.*, Cat. Phal. I 92 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 71, t. 10, ff. K [♂ ♀] (1913).[4] — Upper Tenasserim.[2, 4] Yunnan.[1-4] ——— Hongkong.[3, 4]

huebneri, Boisduval.

Syntomis huebneri, Boisd., Mon. Zyg. p. 127, t. 8, f. 4 (1829).[1]	Bombay.[10, 11]
Syntomis marsdeni, Mo., Cat. Lep. E. I. Co. Mus. II 323 (1859)[2]; Mo., P.Z.S. 1859. 157, t. 60, f. 3 (1859).[3]	Throughout India.[12] Dhamai (E. Bengal.)
Syntomis xanthomela, Wlk., J. Linn. Soc. III 184 (1860).[4]	
Naclia eingulata, Wlgn., Wien. Ent. Mon. IV 39 (1860).[5]	———
Buthisia sangaris, Wlgn., Wien. Ent. Mon. VII 139 (1863).[6]	Singapore.[4, 7, 10, 11]
Syntomis contermina, Wlk., Cat. XXXI 78 (1864).[7]	
Hydrusa pyrrhodera, Meyr., Proc. Linn. Soc. N.S.W. (2) I 777 (1886).[8]	Java.[2, 6, 10, 11]
Syntomis frustulenta, Swinh., Cat. Het. Mus. Oxon. I 44 (1892).[9]	Manila.[5]
Syntomis huebneri, Hmpsn., F. I. Moths I 219 (1892) [redescr.][10]; *id.*, Cat. Phal. I 69 (1898) [redescr.][11]; Seitz, Lep. Indo-Austr. II 74 (1913).[12]	N. Australia.[8, 9, 10, 11]

hydatina, Butler.

Syntomis hydatina, Butl., J. Linn. Soc. XII 346 (1876)[1]; Hmpsn., F. I. Moths I 210-211 (1892) [redescr.] [*Syntomoides*][2]; *id.*, Cat. Phal. I 106; t. 4, f. 14 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 68, t. 10, f. G (1912).[4]	Sikkim.[6] Calcutta.[1-3] Central India.[2-4]

insueta, Swinhoe.

Syntomis insueta, Swinh., Cat. Het. Mus. Oxon. I 41, t. 2, f. 16 (1892)[1]; Hmpsn., F. I. Moths I 213 (1892) ♀[2]; *id.*, Cat. Phal. I 105 (1898) ♀[3]; Seitz, Lep. Indo-Austr. II 69 (1913).[4]	Barrackpur.[1, 4]

khasiana, Butler.

Syntomis khasiana, Butl., J. Linn. Soc. XII 345-346 (1876)[1]; Hmpsn., F. I. Moths I 214 (1892) [redescr.][2]; *id.*, Cat. Phal. I 103, t. 4, f. 9 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 69, t. 10, f. H (1913)[4]; Swinh., T.E.S. 1895. 30 (1895).[5]

Khasis.[1-4] Shillong.[5]

lucina, Butler.

Syntomis lucina, Butl., J. Linn. Soc. XII 345 (1876)[1]; Hmpsn., F. I. Moths I 214 (1892) [redescr.][2]; *id.*, Cat. Phal. I 104, t. 4, f. 10 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 69, t. 10, f. H. (1913).[4]

Khatmandu. Nepal.[1-4] Sikkim.[2-4] Calcutta.[3-4] [? error loci].

luteifascia, Hampson.

Syntomis luteifascia, Hmpsn., F. I. Moths I 218-219 (1892) ♂[1]; *id.*, Cat. Phal. I 97-98, t. 3, f. 25 (1898) ♂[2]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. I (1913).[3]

Nagas.[1-3]

madurensis, Hampson.

Syntomis madurensis, Hmpsn., A.M.N.H. (7) VIII 166 (1901) ♂[1] *id.*, B. J. XVI 151, t. D, f. 27 (1904) [redescr.][2]; Seitz, Lep. Indo-Austr. II 69, t. 9, f. F (1913)[3]; Hmpsn., Cat. Phal. Suppl. I 15, t. 1, f. 23 (1914) ♂ [*Amata*].[4]

Kodaikanal.[4] Madura District.[1-4]

masoni, Moore.

Syntomis masoni, Mo., P.Z.S. 1878. 845, t. 53, f. 4 (1878)[1]; Hmpsn., F. I. Moths I 215-216 (1892) [redescr.][2]; *id.*, Cat. Phal. I 95-96 (1898) [redescr.][3]; Seitz, Lep. Pal. II 39, t. 9, f. F (1910)[4]; *id.*, Lep. Indo-Austr. II 71 (1913).[5]

Upper Tenasserim.[1, 2, 3, 5] Tibet.[2, 4] Moupin.[3, 6]

Syntomis davidi, Poujade, Bull. S.E.Fr. (6) IV 136 (1885)[6] ; Hmpsn., F. I. Moths I 218 (1892)[7] ; Leech, T.E.S. 1898. 323 (1898).[8]

W. China.[4], [8]
S. China.[5]
Central China.[8]

melas, Walker.

Syntomis melas, Wlk., Cat. I 133-134 (1854)[1] ; Butl., Ill. Het. I 17, t. 6, f. 10 (1877)[2] ; Seitz, Lep. Indo-Austr. II 73, t. 9, f. G (1913).[3]

Nepal.[1], [3], [6], [8]
Sikkim.[6], [8]
Assam.[3]
Khasis.[6], [8]

Syntomis andersoni, Mo., P.Z.S. 1871. 244, t. 18, f. 1 (1871)[4]; *id.*, Anderson's Researches in W. Yunnan, p. 926, t. 81, f. 4 (1878).[5]

Shillong.[6], [7]
Chin Hills.[8]

Syntomis melæna, Hmpsn., F. I. Moths I 216 (1892) [emend. pro *melas*] [redescr.][6] ; Swinh., T.E.S. 1895. 31 (1895)[7] ; Hmpsn., Cat. Phal. I 96 (1898) [redescr.].[8]

Yunnan.[3], [4], [5], [8]

minor, Warren.

Syntomis minor, Warr., P.Z.S. 1888, 294 (1888)[1] ; Hmpsn., F. I. Moths I 213 (1892) [redescr.][2] ; *id.*, Cat. Phal. I 107, t. 4, f. 15 (1898) [redescr.][3] ; Seitz, Lep. Indo-Austr. II 69, t. 10, f. H (1913).[4]

Kala Pani (Punjab).[1], [3]
Abbottabad.

muirheadi, Felder.

Syntomis muirheadi, Feld., Wien. Ent. Mon. VI 37 (1862)[1] ; Leech, T.E.S. 1898. 322 (1898)[2] ; Seitz, Lep. Pal. II 40, t. 9. f. G (1910).[3]

Chia-ting-fu [E.Tibet border.][2]

Kwei-chow Prov.[2], [4]
S. China.[3]

Syntomis aucta, Leech, T.E.S. 1898. 323 (1898).[4]

Central China.[2]

Syntomis muirheadi ab. *aucta*, Seitz, Lep. Pal. II 40 (1910).[5]

N. E. China.[2]

newara, Moore.

Syntomis newara, Mo., Lep. Atk. p. 12 (1879)[1]; Hmpsn., F. I. Moths I 217 (1892) ♀ [2]; *id.*, Cat. Phal. I 94, t. 3, f. 18 (1898) ♀ [3]; Seitz, Lep. Indo-Austr. II 70, t. 10 f. i (1913).[4]	Darjiling.[1] Sikkim.[2-4] Bhutan.[2-4]

ochreipuncta, Hampson.

Syntomis ochreipuncta, Hmpsn., F. I. Moths I 222 (1892) ♂ [1]; *id.*, Cat. Phal. I 64, t. 2, f. 17 (1898) ♂ [2]; Seitz, Lep. Indo-Austr. II 74, t. 11, f. B (1913).[3]	Ganjam.[1-3]

owstoni, Rothschild.

Amata owstoni, Roths., Novit. Zool. XVIII 155 (1911)[1]; *id.*, Novit. Zool. XIX 377, t. 5, f. 6 (1912)[2]; Seitz, Lep. Indo-Austr. II 71 (1913) [*Syntomis*].[3]	Yunnan.[4] ——— Annam.[3] Hainan.[1-3]
Amata oustoni, Hmpsn., Cat. Phal. Suppl. I 36 (1914) ♂.[4]	

pasca, Leech.

Syntomis pascus, Leech, T.E.S. 1889. 124, t. 9, f. 1 (1889)[1]; *id.*, T.E.S. 1898. 320-321 (1898).[2]	Tibet.[1] Moupin.[2, 3]
Syntomis pasca, Hmpsn., Cat. Phal. I 94-95 (1898) [redescr.][3]; Seitz, Lep. Pal. II 40, t. 9, f. H (1910).[4]	——— W. China.[2, 4] Central China.[2, 4] Kiukiang.[1, 3]

passalis, Fabricius.

Zygæna cerbera [*nec* Linn.], Sulzer, Gesch. Ins. t. 20, f. 8 (1776).[1]	Assam.[10, 11] Calcutta.[11]
Zygæna passalis, Fabr., Spec. Ins. II 159 (1781).[2]	Pusa.
Sphinx crcusa [*nec* Linn.], Cram., Pap. Exot. III, t. 248, f. F (1788)[3]; Boisd., Mon. Zyg. p. 121, t. 7, f. 7 (1829) [*Syntomis*][4]	Kirkee. Nagpur. Mysore City.

Syntomis latreillii, Boisd., Mon. Zyg. p. 117, t. 7, f. 5 (1829).[5]	Bangalore.[11]
Syntomis montana, Butl., J. Linn. Soc. XII 349 (1876)[6]; Mo., Lep. Ceylon II 36, t. 94, f. 5 (1882).[7]	Coimbatore,[11] Tanjore. Travancore.[14]
Syntomis passalis, Mo., Lep. Ceylon II 35 (1882).[8]	Ceylon.[7, 8, 11]
Zygæna passalis, Kirby, Cat. Het. I 92-93 (1892).[9]	
Syntomis passalis, Hmpsn., F. I. Moths I 219-220 (1892) [redescr.][10]; *id.*, Cat. Phal. I 86 (1898) [redescr.][11]; *id.*, B. J. XX 95 (1910)[12]; Seitz, Lep. Indo-Austr. II 72, t. 11, ff. A [♂♀] (1913)[13]; Fletcher, Proc. Third Entl. Meeting I 52 (1919).[14]	Colombo.[8, 13] Kandy.[6, 13, 15] Peradeniya.
Syntomis passalis ab. *montana*, Seitz., Lep. Indo-Austr. II 72, t. 11, f. A [♂] (1913).[15]	
Larva descr.[12] , on *Ipomœa batatas*.[14]	

pectoralis, Walker.

Syntomis pectoralis, Wlk., Cat. I 133 (1854)[1]; Butl., Ill. Het. I 17, t. 6, f. 5 (1877)[2]; Hmpsn., F. I. Moths I 224 (1892) [redescr.][3]; *id.*, Cat. Phal. I 62-63 (1898) ♀[4]; Seitz, Lep. Indo-Austr. II 73, t. 9, f. E (1913).[5]	Burma.[5] E. Pegu.[2] Moulmein.[1, 3], Dawna Hills.[4] Andamans.[4]
Amata pectoralis, Hmpsn., Cat. Phal. Suppl. I 383 (1914).[6]	————
Hydrusa basirufa, Swinh., Cat. Het. Mus. Oxon. I 51 (1892).[7]	Cambodia.[3-5]
Eressa rubribasis, Joannis, Bull. S. E. France 1912. 287 (1912).[8]	Cochin China.[8]

pentazonata, Hampson.

Syntomis pentazonata, Hmpsn., Cat. Phal. I 99, t 3, f. 27 (1898)[1]; *id.*, B. J. XIII 50 (1900) [redescr.][2]; Seitz, Lep. Indo-Austr. II 72, t. 10, f. L (1913).[3]	N. Chin. Hills.[1, 3]

perixanthia, Hampson.

Syntomis perixanthia, Leech, Entom. XXXI 152 (1898) [non descr.][1]; Hmpsn., Cat. Phal. I 97, t. 3, f. 17 (1898)[2]; Leech, T.E.S. 1898. 321 (1898)[3]; Seitz, Lep. Indo-Austr. II 70 (1913).[4]	Moupin.[2, 3] Ni-tou.[6] Wa-shan.[2, 3] Omei-shan.[2, 3, 6]
Syntomis persimilis, Leech, Entom. XXXI 152 (1898)[5]; Leech, T.E.S. 1898. 322 (1898) ♀.[6]	W. China.[3, 10, 11]
Syntomis perixanthia persimilis, Seitz, Lep. Pal. II 39 (1910).[7]	Formosa.[2, 3, 4, 10]
Amata perixanthia sinensis, Roths., Novit. Zool. XVII 435 (1910)[8]; *id.*, Novit. Zool. XIX 375, t. 3, f. 17 (1912)[9]; Seitz, Lep. Pal. II 39, 445, t. 9, f. F (1910) [*Syntomis*][10]; Hmpsn., Cat. Phal. Suppl. I 36 (1914).[11]	

phœnicozona, Hampson.

Syntomis phœnicozona. Hmpsn., B. J. XI 284 (1897) ♂[1]; *id.*, Cat. Phal. I 63, t. 2, f. 13 (1898) ♂[2]; Seitz, Lep. Indo-Austr. II, 73, t. 11 f. C (1913).[3]	Andamans.[1-3]

quadrifascia, Hampson.

Syntomis quadrifascia, Hmpsn., F. I. Moths I 215 (1892) ♂[1]; *id.*, Cat. Phal. I 100, t. 4 f. 2 (1898) ♂[2]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. K (1913).[3]	Bernardmyo (Burma).[1-3]

rubrozonata, Poujade.

Syntomis rubrozonata, Poujade, Bull. S. E. France (6) VI 117 (1886)[1]; Hmpsn., Cat. Phal. I 85, t. 3 f. 7 (1898) [redescr.][2]; Leech, T. E. S. 1898, 324 (1898)[3]; Seitz, Lep. Pal. II 40, t. 9, f. F (1910).[4]	Moupin.[1, 3, 5], Omei-shan.[2, 3, 7] Tibet.[4]

Syntomis rubrizonata, Kirby, Cat. Het. I 93 (1892).[5]

Syntomis leucoma, Leech, Entom., XXXI, 154 (1898)[6]; *id.*, T. E. S. 1898, 324 (1898).[7] — W. China.[4]

serrata, Hampson.

Syntomis serrata, Hmpsn., F. I. Moths I 221 (1892)[1]; *id.*, Cat. Phal. I 70, t. 2, f. 24 (1898)[2]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. i (1913).[3] — Khasis.[1-4]

Syntomis grotei [*nec* Moore], Swinh., T. E. S. 1895, 31 (1895).[4]

sladeni, Moore.

Syntomis sladeni, Mo., P.Z.S. 1871. 245, t. 18, f. 5 (1871)[1]; *id.*, Anderson's Researches in W. Yunnan, p. 927, t. 81, f. 8 (1878)[2]; Hmpsn., F. I. Moths I 217 (1892) [redescr.][3]; *id.*, Cat. Phal. I 91 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 71, t. 10, f. L (1913).[5]

Yunnan.[1-5]
E. Pegu.[3-5]
Bernardmyo.[3, 4]
Chin Hills.[4]
Upper Tenasserim.[4, 5]

sperbius, Fabricius.

Zygæna sperbius, Fabr., Mantissa II 103 (1787).[1]

Syntomis atkinsoni, Mo., P.Z.S. 1871, 245, t. 18, f. 2 (1871)[2]; Butl., J. Linn. Soc. XII 347 (1876)[3]; Anderson's Researches in W. Yunnan, p. 927, t. 81, f. 5 (1878)[4]; Hmpsn., F. I. Moths I 212 (1892) [redescr.].[5]

Syntomis sperbius, Hmpsn., Cat. Phal. I 106 (1898) [redescr.][6]; *id.*, B. J. XIII 51 (1900)[7]; Seitz, Lep. Indo-Austr. II 68-69, t. 10, ff. G [♂], H [♀] (1912).[8]

Pusa.
Sikkim.[6, 8]
Margherita.
Cachar.[6]
Shillong.
Upper Burma.[6]
Lower Burma.[6]
Moulmein.[2, 6]
Tenasserim.[6]
Yunnan.[4, 5]
———
Tonkin.[6],
Hainan.[8]
China.[1]

stellaris, Snellen.

Syntomis stellaris, Snell., Deuts. Ent. Zeit., Lep. VIII 130 (1895)[1]; Hmpsn., Cat. Phal. I 112, t. 4, f. 22 (1898) ♂.[2]	Tenasserim ? ——— Siam.[2] Perak.[2] Sumatra.[2] Borneo.[2]

submarginalis, Walker.

Syntomis submarginalis, Wlk., Char. Undescr. Het. p. 89 (1869)[1]; Hmpsn., F. I. Moths I 218 (1892) [redescr.][2]; *id.*, Cat. Phal. I 91, t. 3 f. 15 (1898) [redescr.][3]; Swinh., T.E.S. 1898. 31 (1898)[4]; Seitz, Lep. Indo-Austr. II 71, t. 10, f. i (1913).[5]	Sikkim.[2, 3, 5] Khasis.[3, 4] Toungoo (Burma).[3] Bernardmyo.[2, 3]

thoracica, Moore.

Phacusa thoracica, Mo., A.M.N.H. (4) XX 343 (1877)[1]; *id.*, Lep. Ceylon II 38, t. 95, f. 7 (1882).[2]	Ceylon [1-4] Kandy. Peradeniya.[5]
Syntomis thoracica, Hmpsn., F. I. Moths I 224 (1892) [redescr.][3]; *id.*, Cat. Phal. I 63, f. 32 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 73, t. 11, f. B (1913).[5]	

tigrina, Walker.

Syntomis tigrina, Wlk., Cat. XXXI, 71 (1864) ♀.[1]	
Trichæta tigrina, Hmpsn., Cat. Phal. I 54, f. 26 (1898) ♀[2], *id.*, B. J. XIII 46 (1900).[3]	——— Cambodia.[1, 2]
Amata tigrina, Hmpsn., Cat. Phal. Suppl. I 29 (1914).[4]	

[NOTE.—This may be opposite sex of *elwesi* (q. v.)].

unifascia, Hampson.

Syntomis unifascia, Hmpsn., F. I. Moths I 212 (1892) ♀[1]; *id.*, Cat. Phal. I 112, t. 4, f. 20 (1893) ♀[2]; Seitz, Lep. Indo-Austr. II 79, t. 11, f. H (1913).[3] — Sikkim.[1, 3]

verecunda, Swinhoe.

Syntomis verecunda, Swinh., A.M.N.H. (7) IX 78 (1902) ♂[1]; Seitz, Lep. Indo-Austr. II 71 (1913).[2] — Jaintia Hills.[1, 2]

vitrea, Walker.

Syntomis vitrea, Wlk., Cat. VII 1594 (1856)[1]; Hmpsn., F. I. Moths I 218 (1892) ♀[2]; *id.*, Cat. Phal. I 90-91, t. 3, f. 14 (1898) ♂♀[3]; Seitz, Lep. Indo-Austr. II 70, t. 10, f. K (1913).[4] — Assam.[3, 5] Sylhet.[1, 2] Chin Hills.[3, 6, 7]

Syntomis austeni, Mo., P.Z.S. 1879. 389 (1879).[5]

Syntomis polyzonata, Hmpsn., T.E.S. 1895. 283 (1895)♀[6]; *id.*, F. I. Moths IV 466 (1896) ♀.[7]

wimberleyi, Swinhoe.

Syntomis wimberleyi, Swinh., P.Z.S. 1889. 400, t. 43, f. 11 (1889)[1]; Hmpsn., F. I. Moths I 219 (1892) ♀[2]; Hmpsn., Cat. Phal. I 88-89 (1898) ♂♀[3]; Seitz, Lep. Indo-Austr. II 78, t. 11, f. F (1913).[4] — Andamans.[1-4]

xanthograpta, Hampson.

Amata xanthograpta, Hmpsn., Cat. Phal. Suppl. I 32, t. 3, f. 6 (1914) ♂.[1] — Kashmir.[1]

yunnanensis, Rothschild.

Amata yunnanensis, Roths., Novit. Zool. XVIII 155 (1911) ♂ [1]; *id.*, Novit. Zool. XIX 377, t. 5, f. 25 (1912)[2]; Hmpsn., Cat. Phal. Suppl. I 35 (1914) ♂ [3]; Seitz, Lep. Indo-Austr. II 70 (1913) [*Syntomis*].[4] — Yunnan.[1-4]

6. Eressa, Walker (1854).

Eressa, Wlk., Cat. I 149 (1854); Hmpsn., Cat. Phal. I 115, f. 38 (1898) [redescr.]; *id.*, B. J. XIII 223 (1901) [key to spp.]; Seitz, Lep. Pal. II 445 (1912); *id.*, Lep. Indo-Austr. II 82 (1913) [redescr.]. Type: *confinis*, Wlk.

Ctenandra, Felder, Seitz. Akad. Wien. LXIII 38 (1861). Type: *naclioides*, Felder [Amboina].

Trianeura, Butler, J. Linn. Soc., XII 353-354, t. 28, f. 3 (1876). Type: *subaurata*, Wlk.

Chloromeles, Meyr., Proc. Linn. Soc. N.S.W. (2) I 785 (1886). Type: *geographica*, Meyr. [Queensland]

Eressades, Beth.-Baker, A.M.N.H. (8) VII 531 (1911). Type: *flavipunctata*, B.-B. [N. Angola.]

affinis, Moore.

Eressa affinis, Mo., P.Z.S., 1877. 596, t. 59, f. 3 (1877)[1]; Hmpsn., F. I. Moths I 223 (1892) [*partim*; *Syntomis*] [2]; *id.*, Cat. Phal. I 117 (1898) ♂ ♀ [3]; Seitz, Lep. Indo-Austr. II 83 (1913).[4]
Eressa politula, Swinh., Cat. Het. Mus. Oxon. I 52 (1892).[5] — Andamans.[1 4]

annosa, Walker.

Syntomis annosa, Wlk., J. Linn. Soc. III 183 (1859).[1] — Dawna Hills.[5]

Syntomis marcescens, Felder, Reise Novara, Lep. IV, t. 102, f. 12 (1875).[2]	Malay Penins.[7]
Syntomis? lasara, Pagenst., Jahrb. Nass. Verh. XXXVIII 13, t. 2, f. 8 (1885).[3]	Singapore.[1, 4, 5]
	Java.[2, 5, 7]
Syntomis annosa, Hmpsn.,F. I. Moths IV 466 (1896)♀[4]	Nias.[3]
Eressa annosa, Hmpsou, Cat. Phal. I 120 (1898) ♂♀[5]; *id.*, B. J. XIII 224 (1900) [synon.][6]; Seitz, Lep. Indo-Austr. II 82, t. 12, f. B (1913).[7]	

aperiens, Walker.

Syntomis apcriens, Wlk., Cat. XXXI 68 (1864).[1]	Khasis.[7 10]
Trianeura moorei, Butl., J. Linn. Soc. XII 354 (1876).[2]	
Syntomis lydia, Swinh., T.E.S. 1891. 134, t. 8, f. 7 (1891) [aberr.].[3]	Bombay.[2 5]
Syntomis aperiens, Hmpsn., F. I. Moths I 221-222 (1892) ♂ ♀.[4]	Nilgiris.[1, 3, 4, 5]
Eressa apericns, Hmpsn., Cat. Phal. I 119-120 (1898) ♂ ♀[5]; Seitz, Lep. Indo-Austr. II 83 (1913).[6]	Ceylon.[6]
Eressa aperiens khasiana, Roths., Novit. Zool. XVII 438 (1910)[7]; *id.*, Novit. Zool. XIX 376, t. 4, f. 8 (1912)[8]; Hmpsn., Cat. Phal. Suppl. I 48 (1914)[9]; Seitz, Lep. Indo-Austr. II 83, t. 12, f. C (1913).[10]	Sumatra.[6]

confinis, Walker.

Glaucopis confinis, Wlk., Cat. I 149-150 (1854)[1]; *id.*, Cat. VII 1592 (1856) [*Syntomis*].[2]	Simla.[8]
	Kangra.[5, 3]
	Kulu.[6]
Eressa confinis, Mo., Lep. Ceylon II 36, t. 95, f. 6 (1882).[3]	Almora.[6]

Eressa musa, Swinh., P.Z.S. 1885. 290, t. 20, f. 1 (1885)[4]; Hmpsn., F. I. Moths I 222-223 (1892) [redescr.] [*Syntomis*].[5]	Sikkim.[8]
Syntomis confinis, Hmpsn., F. I. Moths I 223 (1892) ♂.[6]	Bhutan.[8]
	Shillong.
Eressa confinis, Swinh., T.E.S. 1895. 32 (1895)[7]; Hmpsn., Cat. Phal. I 116, f. 38 (1898) [redescr.][8]; Seitz. Lep. Indo-Austr. II 83 (1913)[9]; Hmpsn., Cat. Phal. Suppl. I 46-47 (1914) [Synon.].[10]	Cherrapunji.[7]
	Jaintias.[11]
	Calcutta.[8]
	Hoshangabad.
	Bombay.[4, 5, 8]
	Belgaum.[15]
Eressa catoria, Swinh., A.M.N.H. (7) VI 305 (1900).[11] ♂.	Nilgiris.[5, 8]
Syntomoides finitima, Wileman, Entom. XLIII 220 (1910).[12]	S. India.[17]
Eressa confinis malaccensis, Roths., Novit. Zool. XVII 437 (1910)[13]; *id.*, Novit. Zool. XIX 376, t. 4, f. 6 (1912).[14]	Ceylon.[3, 5, 8, 9]
	Burma.[5, 8, 9]
	Myitkyina (U. Burma.)
Eressa confinis intensa, Roths., Novit. Zool. XVII 438 (1910)[15]; *id.*, Novit. Zool, XIX 376, t. 4, f. 7 (1912)[16]; Seitz, Lep. Indo-Austr. II 83, t. 12, f. B (1913).[17]	Penang.[13]
	Tonkin.[13]
	Formosa.[12]

discinota, Moore.

Syntomis discinota, Mo., Lep. Atk. p. 13 (1879)[1]; Hmpsn., F. I. Moths I 216-217 (1892) ♂[2]; Swinh., T.E.S. 1895. 31 (1895).[3]	Khasis.[1-5]
Eressa discinota, Hmpsn., Cat. Phal. I 123, f. 44 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 84, t. 9, f. H (1913).[5]	Tenasserim.[2, 4]

eressoides, Hampson.

Syntomis cressoides, Hmpsn., F. I. Moths I 220 (1892) ♀.[1]	Momeit (Burma).[1, 2]

Eressa eressoides, Hmpsn., Cat. Phal. I 118, t. 4, f. 26 (1898) ♀[2]; *id.*, B. J. XIII 223 (1900)[3]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. B (1913).[4]

erythrosoma, Hampson.

Syntomis erythrosoma, Hmpsn., F. I. Moths I 223-224 (1892) ♂.[1] — Matelei (Burma).[1, 2]

Eressa erythrosoma, Hmpsn., Cat. Phal. I 119, t. 5, f. 23 (1898) ♂[2]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. C (1913).[3]

ichneumoniformis, Rothschild.

Eressa ichneumoniformis, Roths., Novit. Zool. XVII 438 (1910) ♂[1]; *id.*, Nov. Zool. XIX 376, t. 4, f. 10 (1912)[2]; Hmpsn., Cat. Phal. Suppl. I 48 (1914) ♂[3]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. C (1913).[4] — Khasis.[1, 3, 4]

lepcha, Moore.

Syntomis lepcha, Mo., Lep. Atk. p. 12 (1879)[1]; Hmpsn., F. I. Moths I 217 (1892)[2]; *id.*, Cat. Phal. I 121, t. 5, f. 7 (1898) [redescr.][3]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. C (1913).[4] — Darjiling.[1] Sikkim.[2-4]

microchilus, Hampson.

Syntomoides microchilus, Hmpsn., F. I. Moths I 212 (1892).[1] — Tenasserim Valley.[1, 3]

Syntomis plumalis, Hmpsn., F. I. Moths IV 466, (1896) ♂.[2] — Toungoo (Burma).[2, 3]

Eressa microchilus, Hmpsn., Cat. Phal. I 119, t. 5, f. 5 (1898) ♂[3]; *id.*, B. J. XIII 224 (1900) [synon.][4]; Seitz, Lep. Indo-austr. II 84, t. 12, f. C. (1913).[5]

multigutta, Walker.

Syntomis multigutta, Wlk., Cat. I 134 (1854)[1]; Butl., Ill. Het. I 19, t. 7, f. 3 (1877) [*Hydrusa*][2]; Hmpsn., F. I. Moths I 220 (1892) [redescr.][3]; *id.*, Cat. Phal. I 120-121, f. 39 (1898) [redescr.][4]; *id.*, B. J. XIII 224 (1900) [synon.][5]; Leech, T.E.S. 1898. 324 (1898)[6]; Hmpsn., P.Z.S. 1906. 486 (1906)[7]; Seitz, Lep. Pal. II 445 (1912)[8]; *id.*, Lep. Indo-Austr. II 84 (1913).[9]*	Nepal.[1, 3, 4] Sikkim.[2, 4, 11] Gantok.[7] Tibet.[8] Moupin.[3, 4, 10, 1] Omei-shan.[12] Wa-shan.[12] Ni-tou.[6] Shillong. Burma.[3, 4]
Syntomis blanchardi, Poujade, Bull. S. E. France (6) IV 136 (1884)[10]; Hmpsn., F. I. Moths I 221 (1892) ♂[11]; Leech, T.E.S. 1898. 323-324 (1898) [variation][12]; Seitz, Lep. Pal. II, t. 9, f. H (1912).[13]	China.[4, 11]

nigra, Hampson.

Syntomis nigra, Hmpsn., F. I. Moths I 221 (1892) ♂[1]; *id.*, Cat. Phal. I 121, t. 5, f. 8 (1898) ♂[2]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. C (1913).[3]	Simla.[1-3]

subaurata, Walker.

Glaucopis subaurata, Wlk., Cat. I 149 (1854).[1]	Ceylon.[1-4]
Eressa subaurata, Mo., Lep. Ceylon II 36-37, t. 95, f. 4 (1882)[2]; Hmpsn., F. I. Moths I 223 (1892 [redescr.] [*Syntomis*][3]; *id.*, Cat. Phal. I 117 (1898) [redescr.][4]; Seitz, Lep. Indo-Austr. II 83, t. 12, f. B (1913).[5]	Kandy.[5] Peradeniya.

vespa, Hampson.

Eressa vespa, Hmpsn., Cat. Phal. I 118, t. 4, f. 27 (1898) ♂♀[1]; *id.*, B. J. XIII 224 (1900)[2]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. C (1913).[3]	Karen Hills.[1, 2]

* Seitz [9] states that there are two forms [subspecies], (1) *multigutta*, occurring in Burma, and (2) *blanchardi*, ranging from Sikkim to Tibet; but *multigutta* was described from Nepal so that this name cannot be applied to the Burmese form which has as yet no name if sufficiently distinct to deserve one. T. B. F.

vespoides, Rothschild.

Eressa vespoides, Roths., Novit. Zool. XVII 438 (1910)[1]; *id.*, Nov. Zool. XIX 376, t. 4, f. 9 (1912)[2]; Hmpsn., Cat. Phal. Suppl. I 51 (1914) ♂[3]; Seitz, Lep. Indo-Austr. II 84, t. 12, f. D (1913)[4]	Khasis.[1-4]

7. Dysauxes, Hübner (1822).

Dysauxes, Hb., Verz. p. 171 (1822); Hmpsn., Cat. Phal. I 132 (1898) [redescr.]; *id.*, B. J. XIII 224 (1900); Seitz, Lep. Pal. II 41 (1910) [redescr.; habits.]. Type: *ancilla*, Linn. [Europe.]

Naclia, Boisd., Index method, p. 60 (1840); Hmpsn., F. I. Moths I 226 (1892). Type: *ancilla*, Linn.

punctata, Fabricius.

Bombyx punctata, Fb., Spec. Ins. II 207 (1781).[1]	Kashmir.[7, 8, 15, 17]
Noctua ancilla var., Esper. Eur. Schmett. p. 51, t. 85, f. 3 (1786).[2]	
Bombyx serva, Hb., Eur. Schmett. III, ff. 115, 126, 127 (1804 ?).[3]	
Bombyx confamula, Hb., Eur. Schmett. III, ff. 226, 227 (1804 ?).[4]	———
Call. serva, Godart, Lep. France IV 381, t. 39, f. 4 (1822).[5]	Europe.[1-7, 8, 14]
Naclia punctata, Mill., Ann. Soc. Linn. Lyon (2) VIII 234, t. 4, ff. 10, 11 (1861) [larva][6]; Hmpsn., F. I. Moths I 226, f. 142 (1892) [redescr.].[7]	W. Asia.[8, 14]
Dysauxes punctata, Hmpsn., Cat. Phal. I 133-134, f. 53 (1898) [redescr.][8]; *id.*, Cat. Phal. Suppl. I 59-60 (1914) [Synon.][9]; *id.*, *t. c.* p. 383 (1914) [larva][10]; *id.*, B. J. XIII 224 (1900)[11]; Kirby, Butt. Moths Europe, p. 97 [12]; Hofmann, Raupen, p. 38, t. 10, f. 12 [*larva* descr.][13]; Seitz, Lep. Pal. II 41-42, t. 9, f. i. (1910).[14]	

Dysauxes punctata kashmirensis, Roths., Novit. Zool. XVII 441 (1910)[15]; *id.*, Nov. Zool. XIX 376, t. 4, f. 11 (1912)[16]; Seitz, Lep. Pal. II 445 (1912).[17]	Srinagar. Achhabal (Kashmir).
Larva descr.[6], [10], [12], [13], [14] on lichens. [6], [10], [12], [13], [14].	

8. Euchromia, Hübner (1820).

Euchromia Hb., Verz. p. 121 (1820); Wlk. Cat. I 201 (1854); Butl., J. Linn. Soc. XII 363, t. 28, f. 20 (1876); *id.*, T.E.S. 1888. 109 (1888); Meyr., Proc. Linn. Soc. N.S.W. (2) I 786 (1886); Kirby, Cat. Het. I 117 (1892); Hmpsn., F. I. Moths I 226, f. 143 (1892) [redescr.]; *id.*, Cat. Phal. I 293 (1898); *id.*, B. J. XIII 224 (1900) [key to spp.]; Seitz, Lep. Indo-Austr. II 85 (1913) [redescr.]. Type: *guineensis*, Fabr. 1775 (=*sperchius*, Cram. 1777) [W. Africa.]

Phalanna, Wlk. Cat. I 218 (1854). Type: *polymena*, Linn.

Hira, Wlk., Cat I 220 (1854). Type: *guineensis*, Fb.

elegantissima, Wallengren.

Sphinx polymena [*nec* Linn.], Drury, Ill. Exot. Ent. I, t. 26, f. 1 (1773).[1]	Moulmein.[6], [11], [19]
Charida elegantissima, Wlgn., Eugenie's Resa, p. 360 (1861).[2]	Tenasserim.[11] "N. India."[3]
Euchromia orientalis, Butl., J. Linn. Soc. XII 364 (1876)[3]; *id.*, T.E.S. 1888. 114, t. 4, f. 6 (1888)[4]; Hmpsn., F. I. Moths I 227-228 (1892).[5]	[? error loci.] ———— Siam.[9], [11], [20]
Euchromia fraterna, Butl., J. Linn. Soc. XII 364 (1876).[6]	China.[1] Formosa.[11], [18]
Euchromia laura, Butl., J. Linn. Soc. XII 364 (1876)[7]; *id.*, T.E.S. 1888. 114, t. 4, f. 8 (1888).[8]	Hainan.[17] Philippines.[11], [13], [21]
Euchromia siamensis, Butl., J. Linn. Soc. XII 365 (1876).[9]	Celebes.[22]
Euchromia formosana, Butl., T.E.S. 1888. 114, t. 4, f. 7 (1888).[10]	

Euchromia polymena orientalis, Hmpsn. Cat. Phal. I 298 (1898)[11]; *id.*, B. J. XIII 224 (1900).[12]

Euchromia elegantissima var. *diffusihelvola*, Schultz, Philipp. Journ. Gen. Sci. 1908, p. 29, t. 1, f. 3 (1908).[13]

Euchromia polymena elegantissima, Hmpsn., Cat. Phal. Suppl. I 197 (1914) [probably a distinct sp.].[14]

Euchromia polymena ab. *laura*, Seitz, Lep. Indo-Austr. II 85, t. 12, f. D (1913).[15]

Euchromia polymena orientalis, Seitz, Lep. Indo-Austr. II 85 (1913).[16]

Euchromia polymena ab. *kainana*, Seitz, Lep. Indo-Austr. II 85 (1913).[17]

Euchromia polymena formosana, Seitz, *l.c.*[18]

Euchromia polymena fraterna, Seitz, *l. c.*[19]

Euchromia polymena siamensis, Seitz, *l.c.*[20]

Euchromia polymena elegantissima, Seitz, *l.c.*[21]

Euchromia polymena celebensis, Butl., Seitz, *l.c.*[22]

magna, Swinhoe.

Syntomis magna, Swinh., T.E.S. 1891. 133, t. 8, f. 1 (1891).[1] — N. Kanara.[1-3]

Euchromia magna, Hmpsn. F. I. Moths I 227 (1892)[2]; *id.*, Cat. Phal. I 305 (1898)[3]; Seitz, Lep. Indo-Austr. II 86 (1913).[4] — Trincomali.[2-4]

polymena, Linnæus.

Sphinx polymena, Linn. Syst. Nat. (ed. X) I 494 (1758)[1]; Cram., Pap. Exot. I, t. 13, f. D (1775).[2] — Bombay.[6] Bassein Fort (Bombay).

Euchromia polymena, Wlk., Cat. I 219 (1854)[3]; Mo., Lep. Ceylon II 39, t. 94, ff. 6, 6[a] (1882)[4]; Hmpsn., F. I. Moths I 227, f. 143 (1892)[5]; *id.*, Cat. Phal. I 297-298, f. 137 (1898)[6]; Seitz, Lep. Indo-Austr. II 85, t. 12, f. D (1913)[7]; Fletcher, Spol. Zeylan. V 63 (1907)[8]; *id.*, Rept. Second Entl. Meeting, p. 291 (1917)[9]; *id.*, Rept. Third Entl. Meeting I 52 (1919)[10]; Turner, Proc. R. Soc. Victoria (n.s.) XXXV 29 (1922).[11]	Bengal.[6] Laheria Serai. Dacca. Coimbatore. Nilgiris.[6] Travancore.[9], [10] Ceylon.[6], [8] Peradeniya. Upper Burma.[6] Andamans.[6]
Larva descr. [4], [5] on *Ipomœa batatas* [9],[10] Convolvulaceæ.[4]	Sangir.[6] N. W. Australia.[11]

98 Species.

INDEX TO GENERA.

(NOTE.—The numbers refer to the numbers of the genera, not of the pages.)

Agaphthora (Ceryx).
Amata, 5.
Asinusca (Amata).
Asinutea (Amata).

Buthysia (Amata).

Cacosoma (Psichotoë).
Callitomis, 2.
Ceryx, 3.
Chloromelas (Eressa).
Cœnochromia (Amata).
Ctenandra (Eressa).

Dysauxes, 7.

Eressa, 6.
Eressades (Eressa).
Euchromia, 8.

Hira (Euchromia).
Hydrusa (Amata).

Naclia (Dysauxes).

Phalanna (Euchromia).
Psichotoë, 1.

Syntomis (Amata).
Syntomoides (Ceryx).

Trianeura (Eressa).
Trichæta, 4.

INDEX TO SPECIES.

NOTE.—Names *in italics* are treated as synonyms and should be looked up under the names given in roman letters.

acrospila (confluens), Amata.
aetea, Amata.
affinis, Eressa.
albapex, Amata.
albifrons, Amata.
ancilla var. (punctata), Dysauxes.
andersoni (melas), Amata.
annosa, Eressa.
aperiens, Eressa.
approximata (imaon), Ceryx.
artina (imaon), Ceryx.
assamica, Amata.
atereus (diptera), Ceryx.
atkinsoni (sperbius), Amata.
aucta (muirheadi), Amata
aurea, Amata.
aureola, Amata.
austeni (vitrea), Amata.
baiæa, Amata.
baiæa (actea), Amata.
basirufa, Hydrusa (Amata pectoralis).
berinda, Amata.
bicincta, Amata.
blanchardi (multigutta), Eressa.
brachypecten (diptera), Ceryx.

catoria (confinis), Eressa.
celebensis, Euchromia elegantissima.
cerbera (passalis), Amata.
cherra, Ceryx.
chlorocera, Amata.
cingulata, Naclia, (Amata huebneri).
collaris, Zygæna (Amata cyssea).
compta, Amata.
concurrens, Amata dichotoma.

confamula (punctata), Dysauxes.
confinis, Eressa.
confluens, Amata.
congenita, Amata.
connectens, Amata.
consequa, Amata.
contermina (huebneri), Amata.
creusa (passalis), Amata.
cuprea (cyssea), Amata.
cupreipennis (godarti), Ceryx.
cyssea, Amata
cysseoides (cyssea), Amata.

davidi (masoni), Amata.
diaphana, Amata.
dichotoma, Amata.
diffusihelvola, Euchromia elegantissima.
diptera, Ceryx.
discinota, Eressa.
disrupta (divisa), Amata.
divisa, Amata.
duvauceli, Psichotoë.

elegantissima, Euchromia.
elongata, Amata.
elwesi, Amata.
emma (form osæ), Amata.
era, Amata.
eressoides, Eressa.
erythrosoma, Eressa.
euryzona, Amata.
extensa, Amata.

fervida, Amata.
finitima, *Syntomoides* (Eressa confinis).
flavifrons, Amata.
flaviguttata, Amata.
flavolavata, Amata.
formosæ, Amata.
formosana (elegantissima), Euchromia.
fraterna (elegantissima), Euchromia.
frustulenta (huebneri), Amata.
fusiformis (imaon), Ceryx.
fytchei (godarti), Ceryx.

gelatina, Amata.
georgina (cyssea), Amata.
ginorea, Ceryx.
godarti, Ceryx.
grotei, Amata.
grotei (serrata), Amata.

hainana, Euchromia elegantissima.
huebneri, Amata.
hyalina, Ceryx.
hydatina, Amata.

ichneumoniformis, Eressa.
imaon, Ceryx.
incipiens (diptera), Ceryx.
insueta, Amata.
intensa, Eressa confinis.

kashmirensis, Dysauxes punctata.
khasiana, Amata.
khasiana (assamica), Amata.
khasiana, Eressa aperiens.

lasara (annosa), Amata.
laterimacula (pleurasticta), Ceryx.
latreillii (passalis), Amata.
laura (elegantissima), Euchromia.
leechi (confinens), Amata.
lepcha, Eressa.
leucoma (rubrozonata), Amata.
leucosoma, Callitomis.
libera (godarti), Ceryx.
lucina, Amata.
lucina assamensis (Amata assamica).
luteifascia, Amata.
lydia, *Syntomis* (Eressa aperiens).

madurensis, Amata.
magna, Euchromia.
malaccensis, Eressa confinis.
marcescens (annosa), Amata.
marsdeni (huebneri), Amata.
masoni, Amata.
melæna (melas), Amata.
melas, Amata.
microchilus, Eressa.
minor, Amata.
montana (passalis), Amata.
moorei, *Trianeura* (Eressa aperiens).
mota (imaon), Ceryx.
muirheadi, Amata.
multifasciata, Callitomis.
multigutta, Eressa.
musa (confinis), Eressa.

newara, Amata.
nigra, Eressa.

ochreipuncta, Amata.
œnone (diaphana), Amata.
orientalis (elegantissima), Euchromia.
owstoni, Amata.
oustoni (owstoni), Amata.

pasca, Amata.
passalis, Amata.
pectoralis, Amata.
pentazonata, Amata.
perixanthia, Amata.
persimilis (perixanthia), Amata.
phœnicozona, Amata.
pleurasticta, Ceryx.
plumalis (microchilus), Eressa.
politula (affinis), Eressa.
polymena, Euchromia.
polymena (elegantissima), Euchromia.
polyzonata (vitrea), Amata.
punctata, Dysauxes.
pyrrhodera, Hydrusa (Amata hnebneri).

obsoleta, Amata actea swinhoei.

quadrifascia, Amata.
quisqualis (cherra), Ceryx.

rubribasis, Eressa (Amata pectoralis).
rubrozonata, Amata.

sangaris, Buthisia (Amata hnebneri).
sargania (imaon), Ceryx.
schonerrhi (cyssea), Amata.
semicincta, Ceryx.
serrata, Amata.
serva, Bombyx, (Dysauxes punctata).
siamensis (elegantissima), Euchromia.
simplex, Eressa (Trichæta teneiformis).
sinensis, Amata perixanthia.
sladeni, Amata.
sladeni, (connectens), Amata.
sperbius, Amata.
stellaris, Amata.
subaurata, Eressa.
submarginalis, Amata.
swinhoei, Amata actea.
syntomoides, Callitomis.

teneiformis, Trichæta.
thoracica, Amata.
tigrina, Amata.
tigrina (elwesi), Amata.

unifascia, Amata.

verecunda, Amata.
vespa, Eressa.
vespoides, Eressa.
vitrea, Amata.
vitreata (diaphana) Amata.
volans (hyalina), Ceryx.

wimberleyi, Amata.
xanthograpta, Amata.
xanthomela (huebneri), Amata.

GIPC—M—IV-2-80(a)—10-5-25—350.

SMITHSONIAN INSTITUTION LIBRARIES
3 9088 00239246 2
nhent QL483.I4C3X
pt 8 Catalogue of Indian insects.

CALCUTTA
GOVERNMENT OF INDIA PRESS
8, HASTINGS STREET

SMITHSONIAN INSTITUTION LIBRARIES
3 9088 00239246 2
nhent QL483.I4C3X
pt. 8 Catalogue of Indian insects.

CALCUTTA
GOVERNMENT OF INDIA PRESS
8, HASTINGS STREET